

604233
604233

1

Analytical Approximations
Volume 12

Cecil Hastings, Jr.
James P. Wong, Jr.

P-426 ✓ *Bed*

18 August 1953

Approved for OTS release

6 p

COPY	1	OF	1
HARD COPY	\$. 1.00		
MICROFICHE	\$. 0.50 <i>p</i>		

DDC
AUG 19 1964
DDC-IRA C

The RAND Corporation
SANTA MONICA • CALIFORNIA

Copyright 1953
The RAND Corporation

6-11-53

Analytical Approximation

Bessel Function of Imaginary Argument: To better
than .0003 over $(1, \infty)$,

$$e^{-x}I_1(x) \doteq \frac{x}{\sqrt{4.51 + 9.00x + 3.25x^2 + 6.37x^3}}$$

Cecil Hastings, Jr.
James P. Wong, Jr.
RAND Corporation
Copyright 1953

7-1-53

Analytical Approximation

Mach Number in Terms of Pressure Ratio: To better than .001 over $1 \leq M \leq 3$ the inverse of

$$x = \frac{P}{P_R} = \frac{\left[\left(\frac{2\gamma}{\gamma+1} \right) M^2 - \left(\frac{\gamma-1}{\gamma+1} \right) \right]^{\frac{1}{\gamma-1}}}{\left[\left(\frac{\gamma+1}{2} \right) M^2 \right]^{\frac{\gamma}{\gamma-1}}},$$

where $\gamma = 1.4$, is given by

$$M = \frac{8.47 + 35.53x - 25.81x^2}{1 + 32.60x + 6.44x^2}.$$

Cecil Hastings, Jr.
James P. Wong, Jr.
RAND Corporation
Copyright 1953

7-13-53

Analytical Approximation

Mach Number in Terms of Pressure Ratio: To better than
.0014 over $1 \leq M \leq 3$ the inverse of

$$x = \frac{P_S}{P_R} = \frac{\left[\left(\frac{2\gamma}{\gamma+1} \right) M^2 - \left(\frac{\gamma-1}{\gamma+1} \right) \right]^{\frac{1}{\gamma-1}}}{\left[\left(\frac{\gamma+1}{2} \right) M^2 \right]^{\frac{\gamma}{\gamma-1}}},$$

where $\gamma = 1.4$, is given by

$$M \doteq \frac{8.19 + 29.40x - 24.58x^2}{1 + 30x}.$$

Cecil Hastings, Jr.
James P. Wong, Jr.
RAND Corporation
Copyright 1953

Analytical Approximation

Unnamed Definite Integral: To better than .013 over $(0, \infty)$,

$$N(x) = \frac{30}{\pi^4} \int_0^{\infty} \frac{e^{-\left(\frac{x}{t}\right)^7} t^7 dt}{e^{t^2} - 1}$$

$$\approx \frac{1}{1 + .0945x^6 - .0273x^8 + .0029x^{10}}.$$

Cecil Hastings, Jr.
 James P. Wong, Jr.
 RAND Corporation
 Copyright 1953

8-14-53

Analytical Approximation

Pearson Cosine Transformation: To better than .0014
over (0,1),

$$r(x) = \cos \left(\frac{\pi}{1 + \sqrt{x}} \right) \\ \doteq \frac{-1-7.47x+8.47x^2}{1+11.65x+12.05x^2} .$$

$r(x^{-1}) = -r(x)$ can be used to obtain function
values over (1,∞).

Cecil Hastings, Jr.
James P. Wong, Jr.
RAND Corporation
Copyright 1953